

333.91009759  
F6364k

UNIVERSITY  
OF FLORIDA  
LIBRARIES



---

**VEGETATIVE COMMUNITY  
MAPPING OF THE KISSIMMEE  
RIVER FLOODPLAIN BEFORE  
AND AFTER CANALIZATION**

DIVISION OF STATE PLANNING, TALLAHASSEE, FLORIDA

A REPORT TO THE DIVISION OF STATE PLANNING  
FOR THE SPECIAL PROJECT TO PREVENT THE  
EUTROPHICATION OF LAKE OKEECHOBEE

---



KESSIMMEE RIVER FLOODPLAIN VEGETATION  
AND CATTLE CARRYING  
CAPACITY BEFORE AND AFTER CANALIZATION

Prepared By

Buford C. Pruitt  
and  
Stephen E. Gatewood

This report is one of the research reports commissioned by the Florida Division of State Planning as part of the Special Project to Prevent the Eutropication of Lake Okeechobee. This research and others commissioned by the Department of Environmental Regulation and the Central and Southern Florida Flood Control District form the basis for the Special Project final report and management plans.

State of Florida  
Department of Administration  
Division of State Planning  
Bureau of Comprehensive Planning

January, 1976

333.91009759

F6364k



## Table of Contents

	Page
List of Tables	iii
Introduction	1
Research Procedure	2
Results and Discussion	9
References Cited	16
Appendix - Vegetative Communities of the Kissimmee River Floodplain during Pre-and Post-Channelization Periods.	17



Digitized by the Internet Archive  
in 2013

<http://archive.org/details/kissimmeeriverfl00flor>



## List of Tables

Table		Page
1.	Acreage of the vegetative communities of the Kissimmee River Floodplain in the pre-(1954) and post-(1974) canalization periods.	6
2.	Estimated cattle carrying capacity for the Kissimmee River Floodplain vegetative communities, in pre-(1954) and post-(1974) canalization period.	7



## INTRODUCTION

In December of 1974, the Division of State Planning initiated work to map the vegetative communities of the Kissimmee River Floodplain. Objectives were to: (1) quantify acreage and patterns of various marsh communities in the floodplain prior to canalization, (2) quantify acreage and patterns of remaining marsh, improved pasture and upland vegetation that has invaded the floodplain since canalization, (3) assess vegetative trends in the floodplain due to canalization, and (4) apply vegetative acreages to an estimation of total floodplain cattle carrying capacity under pre- and post-canalization conditions.

Results provide baseline information for understanding the natural floodplain ecosystem and quantitative data to evaluate the various reflooding alternatives. Acreage of the various floodplain vegetative communities before and after canalization are multiplied by the approximate cattle stocking rates for each vegetative community to estimate total cattle carrying capacity for the floodplain under both regimes. Acreage figures were used for quantitative evaluations of areas to be reflooded.

## Research Procedure

Natural and cultural habitats of the Kissimmee River Floodplain were outlined on acetate overlays from Agricultural Stabilization and Conservation Service aerial photographs at a scale of 1:12,000, which were before (1952-54) and after (1974) construction of Canal C-38. After ground truthing, blueprinted copies of each habitat were cut out and tabulated by an automatic area meter (Model AAM-5, Hyashi Denko Co., Tokyo, Japan) to derive acreages. Recommended cattle carrying capacities for the various habitats were derived from Soil Conservation Service Range Technicians Guides, the Gatewood and Cornwell (1975) study of the Kissimmee Basin cattle industry and Dr. L.D. White (pers. communication, 1975).

The floodplain area quantified lies south of State Highway 60 and north of State Highway 70, between the original upland vegetative habitats as interpreted from aerial photographs. Sloughs and streams were excluded upstream from their junction with the floodplain. The area north of the old "Government Cut," extending about 7 miles upstream from Lake Okeechobee was excluded as it existed in 1954 and had already experienced changes due to canalization, changing little in land use from 1954 to 1974.

Five generalized vegetative communities were identified from pre-canalization (1954) aerial photos. Two additional habitats were identified from post-canalization (1974) aerial photos (Table 1). These habitats were defined as follows:

Improved pasture - pastureland managed for a few selected species of tame forage grasses after completion of drainage and removal of native vegetation. The most commonly used pasture grasses are bermudagrass (Cynodon dactylon), bahiagrass (Paspalum notatum), pangolagrass (Digitaria decumbens), and their varieties. Pastures are often separated from surrounding habitats through ditching and/or diking practices.

Unimproved pasture - pastureland which contains a mixture of forage and non-forage species, both native and introduced. Some representative taxa are carpetgrass (Axonopus spp.), panic grasses (Panicum spp.), paspalums (Paspalum spp.), bluestems (Andropogon spp.), sedges (Carex spp. and Cyperus spp.) and such forbs as pennywort (Centella asiatica) and dogfennel (Eupatorium spp.) It may contain some shrub species such as salt bush (Baccharis halmifolia) and wax myrtle (Myrica cerifera), usually as scattered individuals or in small patches. Drainage control is minimal.

Shrubby pasture - grassland pasture covered with 15-20 percent or more of shrub species. This habitat occurs where shrub species such as wax

myrtle and salt bush have invaded pastures adjacent to upland habitats and is used as a relative measure of shrub invasion into 1954 pasture.

Marshland - areas including both herbaceous and mixed shrub wetlands and characterized by one or more of the following taxa:

Maidencane	<u>Panicum hemitomon</u>
Cutgrasses	<u>Leersia spp.</u>
Watergrass	<u>Hydrochloa caroliniensis</u>
Sedges	Cyperaceae
Rushes	Juncaceae
Cattail	<u>Typha spp.</u>
Pickereel weed	<u>Pontederia spp.</u>
Arrowhead	<u>Sagittaria spp.</u>
Primrose willow	<u>Ludwigia spp.</u>
Willow	<u>Salix spp.</u>
Buttonbush	<u>Cephalanthus occidentalis</u>
Elderberry	<u>Sambucus canadensis</u>

Shrubland - areas with a relatively dense shrub canopy, approximately 50 percent coverage or greater. This classification includes both wet and dry shrub vegetation types.

Open water and disturbed habitats (ruderal) - areas including the Kissimmee River and its backwaters and tributaries, Canal C-38 ponds, numerous large ditches and dikes, levees, highway and railroad rights-of-way and buildings with their immediate surroundings.

Spoil areas - deposits of spoil dredged in construction of C-38, much of which is barren and supports little or no vegetation. Acreages of grazeable spoil are included in Table, 1, which lists acreage of



floodplain vegetative communities in pre- and post-canalization periods.

The habitats are outlined in Figures 2a and b through 20a and b, and are arranged with corresponding floodplain segments of pre- and post-canalization conditions in Appendix A.

Cattle carrying capacity estimates reported are approximations based on Soil Conservation Service recommended stocking rates for floodplain vegetation associations, expressed as the number of acres required to sustain one cow for a year. These are the best available rates. Each habitat type has a range of values which depends on its condition class, judged as excellent, good, fair or poor. Since determination of conditions class was not possible, Table 2 presents maximum and minimum stocking rates from excellent and poor condition classes, respectively. The computed cattle carrying capacity is the value derived using the median of the two extremes. This would correspond to an "average" condition class being applied to all habitats.

The stocking rates presented are estimates based on the amount of forage produced by the major vegetation associations identified. Each association is composed of various communities within the major grouping. For example, marshland is composed of numerous, intermingled wetland communities with different grazing values. Some are

TABLE I

Acreage of the vegetative communities of the Kissimmee River Floodplain in pre-(1954) and post-(1974) canalization periods.

Habitat	1954			1974			Change	
			% of Total Acres			% of Total Acres	Acreage	%
	Acres			Acres				
1. Pasture								
a. Improved	900	1.4		15,400 <sup>1</sup>	29.6		+14,500	+1,600
b. Unimproved	5,090	9.8		9,020 <sup>1</sup>	17.4		+ 3930	+ 7.7
c. Shrubby	-	-		3,760 <sup>1</sup>	7.2		+ 3,760	-
Total	5,990	11.2		28,180	54.2		+22,190	+ 370
2. Marshland	40,600	78.2		8,840	17.0		-31,760	- 78
3. Shrubland	550	1.1		2,170	4.2		+ 1,615	+ 291
4. Open Water and Disturbed (Ruderal)	4,750	9.1		5,380	10.4		+ 630	+ 13
5. Spoil Areas								
a. In Pasture	-	-		1,300 <sup>1</sup>	2.5		+ 1,300	-
b. Unused	-	-		7,380	14.2		+ 7,380	-
Total	-	-		8,680	16.7		+ 8,680	-
Totals	51,890	-		51,950	-		-	-

<sup>1</sup>Acreage of spoil area pasturage included in above (1(a-c)) pastures.



TABLE 2

Estimated cattle carrying capacity for the Kissimmee River Floodplain vegetative communities, in pre-(1954) and post-(1974) canalization periods.

Habitat	1954			1974		
	Acreage	Acres Per AU	Computed Cattle Carrying Capacity (Number of AU) <sup>1</sup>	Acreage	Acres Per AU	Computed Cattle Carrying Capacity (Number of AU)
1. Pasture						
a. Improved	900	1- 5	300	15,400	1- 5	5,130
b. Unimproved	5,090	5-15	510	9,020	5-15	900
c. Shrubby	- -	- -	- -	3,760	5-15	380
Total	5,990	- -	810	28,180	- -	6,410
2. Marshland	40,600	2-12	5,800	8,840	2-20	800
3. Shrubland	550	5-20	40	2,170	5-20	170
4. Open Water and Disturbed (Ruderal)	4,750	- -	- -	5,380	- -	- -
5. Spoil Areas						
a. In Pasture	- -	- -	- -	1,300 <sup>2</sup>	- -	- -
b. Unused	- -	- -	- -	7,380	- -	- -
Total	- -	- -	- -	8,680	- -	- -
Total	51,890 <sup>3</sup>	- -	6,650	51,950 <sup>3</sup>	- -	7,380 <sup>4</sup>

1. Estimations in this column are median values as described in the text
2. Cattle carrying capacity included in 1(a-c) above
3. Error in acreage values computed at 0.053%
4. Increase of 11%

dominated by preferred forage plants (maidencane, cutgrass), while others by low value plants (cattail, pickerelweed). The range condition class is based on the abundance of preferred forage species for that habitat as follows:

<u>Class</u>	<u>Percent of Present Vegetation Potential of Site</u>
Excellent	76-100
Good	51-75
Fair	26-50
Poor	0-25

Thus, using a median value between excellent and poor stocking rates assumes that high and low forage value areas balance each other out to an "average" condition.

It should be emphasized that all of the area in question has not been used necessarily for cattle production in the past. However, for the purpose of estimating total floodplain carrying capacity it is assumed that all grazeable acreage could be utilized, including land from which cattle are presently excluded by fences, the river, deep marsh water or canals. Values for carrying capacities are estimated from the total acreage of vegetative habitats for the entire floodplain as if it were managed as one production unit.

Also of prime importance is consideration of the effects flood waters have on actual carrying capacity. The pre-canalization condition was characterized by seasonal flooding on the floodplain which affected cattle use. However, this occurred at a time when proper marsh

management calls for resting marsh pastures from grazing. High flood waters provided a natural deferment during the growing season, permitting preferred forage species to maintain vigor and reproduce. Forage production accumulated during limited access for subsequent grazing as flood waters receded. The change in actual carrying capacity due to high water could not be specifically quantified with the information available.

Determination of precise cattle carrying capacities for the floodplain is not possible without a parcel-by-parcel evaluation of the area. It is influenced by: (1) types and amount of forage produced, (2) previous management practices, (3) local variations in soils and corresponding vegetative cover, (4) ground elevation in relation to fluctuating water levels, and other factors (Gatewood and Cornwell, 1975). However, this technique provides an approximation based on the acreage of generalized vegetative communities identified and may be used for comparative purposes. Further quantification of the many interacting factors would permit more exact estimates of cattle carrying capacity, but are beyond the scope of this study.

### Results and Discussion

Profound and obvious changes have occurred to Kissimmee River Floodplain vegetative communities since 1954. Construction of Canal C-38 with subsequent floodplain drainage and shifts in land management techniques have changed vegetative patterns completely. Marshland has been reduced by 78 percent. Pasture has increased by 370 percent, the majority replacing marshland. Spoil areas now cover 8,680 acres of former floodplain habitats.

Open water and ruderal acreages have increased little, yet their character has changed radically. Various aspects of this process of change have been further described by Goodrick and Milleson (1974), Gatewood and Cornwell (1975) and Marshall, et. al. (1972).

The 1954 riverine marshland averaged one mile wide and covered 40,600 acres (63.4 square miles) which supported a number of vegetative communities. Large stands of maidencane and cutgrass were interspersed within the large community which included a diverse group of sedges, forbs, grasses and shrubs. Smaller marshes of similar composition were common in depressions along the floodplain fringes and where numerous creeks and sloughs entered the floodplain. Oxbow lakes and small, shallow ponds were plentiful.

Marshland by 1974 had been reduced to 8,840 acres (13.8 square miles). Most of this was confined to the five pools behind the control structures on C-38. Stabilized water levels in the pools have created deep marshes which differ from those present under fluctuating water levels.\* Cattle prefer not to forage in deeper areas of the pools due to deep water year-round. Subsequently, emergent vegetation is very tall and dense.

---

\*Some pools are now being fluctuated approximately one foot on an annual basis (Goodrick and Milleson, 1974).

Pasture vegetation in 1954 was confined to 5,990 acres of the floodplain's higher fringes. About 85 percent of the pasture was unimproved, with cattle generally grazing on a seasonal basis depending on water levels.

Pasture in 1974 had been increased to 28,180 acres encompassing 54 percent of the floodplain. Improved pasture now occupied almost 30 percent of the floodplain, up from 1.4 percent in 1954. Pasture covered most of the land potentially available for intensive management by cattlemen and replaced large areas of former marshland.

Shrubland in 1954 was generally limited to natural river levees and to occasional willow-buttonbush swamps along floodplain fringes and covered about 550 acres (1.1 percent) of the floodplain. There was very little invasion by upland shrubs into floodplain fringe pastures.

Shrubland had increased 291 percent to 2,170 acres by 1974. Shrubs occurred on agricultural and flood control levees, spoil areas and on some islands formed between the old riverbed and C-38. Shrubs had also invaded a large portion (74 percent) of fringe pastureland.

Open water and ruderal areas occupied 4,750 acres in 1954. Most open water was contained in the Kissimmee River, its oxbow lakes and by-passes, and numerous small ponds scattered throughout the floodplain. Ruderal areas consisted mostly of agricultural levees and railroad or highway rights-of-way.

Open water and ruderal had increased only 13 percent by 1974; however, its character had changed radically. Most open water was confined to the relatively slow moving waters of C-38



with its five impoundments and the larger former river channels. Ruderal areas increased with the addition of many new flood control and agricultural levees.

The spoil banks of dredged material from C-38 covered 8,680 acres of the floodplain. Within these banks were 7,380 acres of bare spoil supporting little or no vegetation. Herbaceous and shrubby vegetation covered the remaining 1,300 acres.

From these changes in vegetative habitats it is estimated that the cattle carrying capacity of the floodplain in 1974 was increased only 11 percent over that which may have been possible under the pre-canalized system. This increase is due primarily to shifts in total acreage of marshland and improved pasture habitats. Marshland supported most of the cattle prior to canalization while the 1974 system carried most of the cattle on improved pasture.

The estimated number of cattle that could be carried on the vegetative habitats existing prior to canalization was 6,650 cows. This was increased to 7,380 under 1974 conditions, yielding average flood plain stocking rates of 7.8 acres/cow and 7.0 acres/cow, respectively.

Shifts in acreages of marshland and pasture had the most significant effect on total floodplain carrying capacity calculations. Marshland in 1954 carried 87 percent of the estimated load with pasture contributing 12 percent. In 1974, the increased acreage of pasture carried 87 percent of the estimated cattle with reduced marshland supporting 11

percent. As was expected, the roles of marshland and pasture for sustaining floodplain cattle have been exchanged as a result of canalization.

The character of grazing resources has changed along with actual acreages. The predrainage (1954) floodplain was characterized by seasonal flooding, which displaced cattle and provided a natural deferment from grazing for marshland habitats. The construction of C-38 water level control combined with the close proximity of remaining marshland to improved pastures established within the floodplain, provided cattle with year-round access to remaining marsh sites. This permits overgrazing of desirable native forages by the high cattle densities present on improved pasture and encourages the growth of plants with low forage value. The primary grazing resource has been changed from one of naturally maintained native forages to one of culturally maintained tame forages.

Overall conditions in the Kissimmee River Floodplain between 1954 and 1974 can be characterized by three major shifts in structure and function: (1) plant communities and land management techniques have been changed from systems requiring low energy inputs to systems requiring high energy inputs, (2) 1974 systems were culturalized totally, while 1954 systems were only partially culturalized, and (3) a floodplain marsh system with naturally maintained high productivity was replaced by a system of pastures with slightly higher, culturally maintained productivity.

The 1954 floodplain ecosystem was one of extensive marshes and grasslands grazed by free roaming cattle. This required very little machinery, low amounts of fossil fuels and little or no fertilizers. By 1974, conditions were such that large amounts of gasoline and fuels, much farm machinery and substantial amounts of fertilizers had been used in developing and maintaining the pasture ecosystem. It can be seen that large amounts of capital are required to sustain the present floodplain system.

The floodplain was only partially culturalized in 1954; that is, very little human influence was used to subsidize natural driving forces. Water levels were governed by basin rainfall and the hydrologic capabilities of the Kissimmee River. Nutrients were deposited by seasonally flooding river waters. Vegetation was controlled by natural water levels, periodic cattle grazing and limited human management. By 1974, as a direct result of construction of C-38, the floodplain had become completely culturalized. No aspect of its functioning escaped human influence. Water levels were dictated by human management policies. Most natural drainage patterns within the floodplain were altered by ditching. Fertilizers were being applied only to improved pastures while remaining areas no longer received nutrient enrichment by flood waters. Vegetation was controlled by manipulated water levels and through intensive management by land owners. Most natural driving forces in operation prior to canalization were no longer effective.



Natural communities, especially marshland, had high potential forage productivity for cattle. Desirable native forage was abundant or dominant in many locations. This natural system was estimated to have only slightly lower overall productivity (in terms of estimated cattle supported) than the 1974 system. While improved pasture was over twice as productive on a per acre basis as marshland, which is evidenced by the ability to carry almost as many cattle on less than half the acreage, the natural system maintained its productivity at far lower cost to man than is required to sustain present culturalized systems.

It thus appears that the increase in total floodplain cattle carrying capacity is estimated to be very low. This small increase may not be significant in light of the high costs incurred in establishing improved pastures and maintaining long term productivity. From a cost-benefit viewpoint, Canal C-38 may have actually had an adverse effect on cattle production in the Kissimmee River Floodplain by requiring substantial cultural subsidies to maintain productivity.

## REFERENCES CITED

- Gatewood, S. E. and G. W. Cornwell. 1975. An analysis of cattle ranching in the Kissimmee River Basin. Florida Department of Administration, Division of State Planning, Bureau of Comprehensive Planning.
- Goodrick, R. L. and J. F. Milleson. 1974. Studies of flood-plain vegetation and water level fluctuation in the Kissimmee River Valley. Central and Southern Florida Flood Control District, March.
- Marshall, A. R., et. al. 1972. The Kissimmee - Okeechobee Basin: A Report to the Cabinet of Florida. Division of Applied Ecology, Center for Urban and Regional Studies, University of Miami, December 12.

## Appendix

Vegetative Communities Maps of the Kissimmee  
River Floodplain during Pre- and Post-Channelization  
Periods.



# **INDEX Map FOR FIGURES 2A THRU 20b**




FIGURE 1





# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post -Channelization Pool A

## POST-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh
- SHR - Shrubby
- RU - Ruderal (old river bed, ditches, canals, etc.)
- SP - Spoil Banks
- Open Water




FIGURE 2b







# VEGETATIVE COMMUNITIES OF THE KISSIMEE RIVER flood plain PRE-Channelization

This map illustrates the vegetative communities of the Kissimmee River flood plain prior to channelization. The river is depicted as a winding, meandering waterway, with its banks and surrounding areas labeled with various codes. The codes are defined in the legend: PAS - Pasture, Im - Improved, Un - Unimproved, MAR - Marsh, RU - Ruderal (levees, spoil, impoundments, etc.), SHR - Shrubby, and Open Water. The map shows a complex network of marshes (MAR) and pastures (PAS) along the river's course. A north arrow is located in the bottom right corner.


**PRE-CONSTRUCTION**

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- MAR - Marsh
- RU - Ruderal (levees, spoil, impoundments, etc.)
- SHR - Shrubby
- Open Water

- PAS - Pasture
  - Im - Improved
  - Un - Unimproved
- MAR - Marsh
- RU - Ruderal (leaves, spoil,  
impoundments, etc.)
- SHR - Shrubby
  - Open Water

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool A

Palmetto  
Flatwoods-  
Pasture



## Post-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh
- SHR - Shrubby
- RU - Ruderal (old river bed, ditches, canals, etc)
- SP - Spoil Banks
- Open Water

Tree & Palmetto  
PAS




FIGURE 4b



Skeeter Slough



FIGURE 5a



Pine Island Slough

PAS - Pasture

Im-Improved

Un - Unimproved

MAR - Marsh

RU - Ruderal (levees, spoil, impoundments, etc.)

SHR - Shrubby

- Open Water

Dense Upland

SHR-  
PAS-  
Upland

SHR-  
PAS-  
Upland

MAR

PAS

MAR

MAR-  
RUD

MAR

MAR

277

MAR

MAR

PAS-

21

Dense  
Trees -  
Palmettos

MAR

Tick Island  
Slough

Dense  
Trees -  
Palmetto

PAS

③

1

MAR

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-CHANNELIZATION







FIGURE 6b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-CHANNELIZATION

## PRE-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- MAR - Marsh
- RU - Ruderal (levees, spoil, impoundments, etc.)
- SHR - Shrubby
- Open Water





# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool B

## POST-CONSTRUCTION

PAS - Pasture

Im - Improved

Un - Unimproved

Shr - Shrubby

MAR - Marsh

SHR - Shrubby

RU - Ruderal (old river bed,  
ditches, canals, etc.)

SP - Spoil Banks

- Open Water

*Trees & Palmettos*





FIGURE 7b



## PRE-CONSTRUCTION

PAS - Pasture  
 Im - Improved  
 Un - Unimproved  
 MAR - Marsh  
 RU - Ruderal (levees, spoil, impoundments, etc.)  
 SHR - Shrubby  
 - Open Water

## VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-CHANNELIZATION

# POST-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh
- SHR - Shrubby
- RU - Ruderal (old river bed, ditches, canals, etc.)
- SP - Spoil Banks
- Open Water

## VEGETATIVE COMMUNITIES OF THE KISSIMEE RIVER FLOOD PLAIN Post-Channelization Pool B





FIGURE 8b



34



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool B




FIGURE 9b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain

## PRE-CHANNELIZATION

### PRE-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- MAR - Marsh
- RU - Ruderal (levees, spoil, impoundments, etc.)
- SHR - Shrubby
- Open Water





FIGURE 10a

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain

## Post-Channelization Pool B



### Post-Construction

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- shr - Shrubby
- MAR - Marsh

SHR - Shrubby


RU - Ruderal (old river bed,  
ditches, canals, etc.)

SP - Spoil Banks  
- Open Water

FIGURE 10b



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER FLOOD PLAIN Pre-Channelization



## Pre-Construction

PAS - Pasture  
Im - Improved  
Un - Unimproved  
MAR - Marsh

RU - Ruderal (levees, spoil, impoundments, etc.)

SHR - Shrubby

- Open Water

FIGURE 11a

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER FLOOD PLAIN Post -Channelization

## Pool B,C




FIGURE 11b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-Channelization

Hydro-  
station  
Stations




FIGURE 12a



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post ~CHANNELIZATION Pool C




FIGURE 12b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-Channelization




FIGURE 13a

# VEGETATIVE COMMUNITIES OF THE kissimmee river flood plain Post-Channelization Pool C





FIGURE 13b



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Pre-Channelization



## PRE-CONSTRUCTION

PAS - Pasture  
Im - Improved  
Un - Unimproved

MAR - Marsh  
RU - Ruderal (levees, spoil, impoundments, etc.)  
SHR - Shrubby  
- Open Water



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post -Channelization Pool C,D




FIGURE 14b

# Pre-Channelization





# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post -Channelization Pool D





FIGURE 15b



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool D





FIGURE 16b



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER FLOOD PLAIN

## Pre-Channelization



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool D





FIGURE 17b



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER FLOOD PLAIN Post -Channelization



# POST-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh
- SHR - Shrubby
- RU - Ruderal (old river bed, ditches, canals, etc.)
- SP - Spoil Banks
- Open Water





FIGURE 18b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-CHANNELIZATION



## POST-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh

SHR - Shrubby

RU - Ruderal (old riverbed, ditches, canals, etc.)

SP - Spoil Banks  
- Open Water



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool E

## POST-CONSTRUCTION

- PAS - Pasture
- Im - Improved
- Un - Unimproved
- Shr - Shrubby
- MAR - Marsh
- SHR - Shrubby
- RU - Ruderal (old riverbed, ditches, canals, etc.)
- SP - Spoil Banks
- Open Water





FIGURE 19b

# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain PRE-CHANNELIZATION



**PRE-CONSTRUCTION**

PAS - Pasture  
Im - Improved  
Un - Unimproved  
MAR - Marsh

RU - Ruderal (levees, spoil, impoundments, etc.)  
SHR - Shrubby  
- Open Water



# VEGETATIVE COMMUNITIES OF THE KISSIMMEE RIVER flood plain Post-Channelization Pool E




FIGURE 20b







333.91007-57

F6364R



## Date Due

Due	Returned	Due	Returned
APR 1 1987			
APR 0 5 1988	APR 0 8 1987		
MAY 2 5 1988			
	MAY 1 8 1988		
	MAY 2 3 1988		
JUN 1 7 1988	MAY 3 1 1988		
	MAR 2 3 1990		
APR 1 3 1990			
MAY 0 4 1990	APR 2 7 1990		
	APR 2 7 1990		
JUL 0 1 1996			
DEC 2 0 1996	DEC 2 0 1996		
APR 0 1 1997	SEP 1 5 1997		

Kissimmee River floodplain UGG  
333.91009759 F6364k



3 1262 01040 7541

